

CSE 250: Math Refresher

Lecture 2

Guest Lecturer: **Aaron Huber**

Sept 1, 2023

Reminders

- AI Quiz due Weds, Sept 6 at 11:59 PM.
 - Your final submission must have a score of 1.0 to pass the class.
 - If you can't submit in autolab, let course staff know ASAP.
- PA 0 due Sun, Sept 10 at 11:59 PM.
 - All you need to do is make sure you have a working environment.
 - If you can't submit in autolab, let course staff know ASAP.
- WA1 to be assigned Monday (Topic: Today's lecture)

Math Refresher

- 1 Summations
- 2 Logarithms
- 3 Limits

Summations

$$\sum_{i=j}^k f(i) = f(j) + f(j+1) + \dots + f(k)$$

Useful Tricks

If c is a constant:

$$\begin{aligned}\sum_{i=j}^k c &= \underbrace{c + \dots + c}_{(k-j+1) \text{ times}} \\ &= (k - j + 1) \cdot c\end{aligned}$$

Useful Tricks

If c is a constant and $f(i)$ is a function of i :

$$\begin{aligned}\sum_{i=j}^k c \cdot f(i) &= c \cdot f(j) + c \cdot f(j+1) + \dots + c \cdot f(k) \\ &= c \cdot (f(j) + f(j+1) + \dots + f(k)) \\ &= c \cdot \sum_{i=j}^k f(i)\end{aligned}$$

Useful Tricks

If $f(i)$ and $g(i)$ are functions of i :

$$\begin{aligned}\sum_{i=j}^k f(i) + g(i) &= (f(j) + g(j)) + \dots + (f(k) + g(k)) \\ &= (f(j) + \dots + f(k)) + (g(j) + \dots + g(k)) \\ &= \left(\sum_{i=j}^k f(i) \right) + \left(\sum_{i=j}^k g(i) \right)\end{aligned}$$

Useful Tricks

If $j < \ell \leq k$:

$$\begin{aligned}\sum_{i=j}^k f(i) &= f(j) + \dots + f(k) \\ &= f(j) + \dots + f(\ell - 1) + f(\ell) \dots + f(k) \\ &= \left(\sum_{i=j}^{\ell-1} f(i) \right) + \left(\sum_{i=\ell}^k f(i) \right)\end{aligned}$$

Useful Tricks

If $j < \ell \leq k$:

$$\left(\sum_{i=j}^k f(i) \right) = \left(\sum_{i=j}^{\ell-1} f(i) \right) + \left(\sum_{i=\ell}^k f(i) \right)$$
$$\left(\sum_{i=j}^k f(i) \right) - \left(\sum_{i=j}^{\ell-1} f(i) \right) = \left(\sum_{i=\ell}^k f(i) \right)$$

Series

$$\sum_{i=1}^k i = \frac{k(k+1)}{2}$$

$$\sum_{i=0}^k 2^i = 2^{k+1} - 1$$

Summary

$$\sum_{i=j}^k c = (k - j + 1) \cdot c \quad (1)$$

$$\sum_{i=j}^k c \cdot f(i) = c \cdot \sum_{i=j}^k f(i) \quad (2)$$

$$\sum_{i=j}^k f(i) + g(i) = \left(\sum_{i=j}^k f(i) \right) + \left(\sum_{i=j}^k g(i) \right) \quad (3)$$


$$\sum_{i=j}^k f(i) = \left(\sum_{i=j}^{\ell-1} f(i) \right) + \left(\sum_{i=\ell}^k f(i) \right) \quad (4)$$

$$\left(\sum_{i=\ell}^k f(i) \right) = \left(\sum_{i=j}^k f(i) \right) - \left(\sum_{i=j}^{\ell-1} f(i) \right) \quad (5)$$

$$\sum_{i=1}^k i = \frac{k(k+1)}{2} \quad (6)$$

$$\sum_{i=0}^k 2^i = 2^{k+1} - 1 \quad (7)$$

Logarithms



Stepping back

$$a \cdot n = \underbrace{a + a + \dots + a}_{n \text{ times}}$$

$$a^n = ? \underbrace{a \cdot a \cdot \dots \cdot a}_{n \text{ times}}$$

Logarithms

$\log_a(b)$ = the number of times you multiply a together to get b

Examples

$$\log_2(32) = 5$$

$$\log_3(27) = 3$$

$$\log_2\left(\frac{1}{8}\right) = -3$$

$$\log_2(2^{10}) = 10$$

$$32 = \underbrace{2 \cdot 2 \cdot 2 \cdot 2 \cdot 2}_{5 \text{ times}}$$

$$27 = \underbrace{3 \cdot 3 \cdot 3}_{3 \text{ times}}$$

$$\frac{1}{8} = 1 \underbrace{\div 2 \div 2 \div 2}_{-3 \text{ times}}$$

$$2^{10} = \underbrace{2 \cdot \dots \cdot 2}_{10 \text{ times}}$$

Logarithm is the Inverse Exponent

$$b^{\log_b(n)} = n = \log_b(b^n)$$

The Product Rule

Let's say $n = a \cdot b$.

How are $\log_2(n)$, $\log_2(a)$, and $\log_2(b)$ related?

$$a = \underbrace{2 \cdot \dots \cdot 2}_{\log_2(a) \text{ times}}$$

$$b = \underbrace{2 \cdot \dots \cdot 2}_{\log_2(b) \text{ times}}$$

$$n = \underbrace{2 \cdot \dots \cdot 2}_{\log_2(n) \text{ times}} = \underbrace{2 \cdot \dots \cdot 2}_{\log_2(a) \text{ times}} \cdot \underbrace{2 \cdot \dots \cdot 2}_{\log_2(b) \text{ times}}$$

$$\log_2(n) = \log_2(ab) = \log_2(a) + \log_2(b)$$

The Exponent Rule

$$\begin{aligned}\log_2(a^n) &= \log_2(\underbrace{a \cdot \dots \cdot a}_{n \text{ times}}) \\ &= \log_2(a) + \log_2(\underbrace{a \cdot \dots \cdot a}_{n-1 \text{ times}}) \\ &= \log_2(a) + \log_2(a) + \log_2(\underbrace{a \cdot \dots \cdot a}_{n-2 \text{ times}}) \\ &= \underbrace{\log_2(a) + \dots + \log_2(a)}_{n \text{ times}} \\ &= n \cdot \log_2(a)\end{aligned}$$

The Division Rule

$$\begin{aligned}\log_2\left(\frac{a}{b}\right) &= \log_2\left(a \cdot \frac{1}{b}\right) \\ &= \log_2(a) + \log_2\left(\frac{1}{b}\right) \\ &= \log_2(a) + \log_2(b^{-1}) \\ &= \log_2(a) + (-1) \cdot \log_2(b) \\ &= \log_2(a) - \log_2(b)\end{aligned}$$

Change of Base

$$b^m = n$$

$$\log_c(b^m) = \log_c(n)$$

$$m \log_c(b) = \log_c(n)$$

$$m = \frac{\log_c(n)}{\log_c(b)}$$

$$\log_b(n) = \frac{\log_c(n)}{\log_c(b)}$$

Summary

Exponent Rule	$\log(n^a) = a \log(n)$
Product Rule	$\log(ab) = \log(a) + \log(b)$
Division Rule	$\log\left(\frac{a}{b}\right) = \log(a) - \log(b)$
Change of Base	$\log_b(n) = \frac{\log_c(n)}{\log_c(b)}$
Inverse	$b^{\log_b(n)} = \log_b(b^n) = n$

For this class, assume base-2 logarithms unless stated otherwise.

Limits

$$\lim_{i \rightarrow c} f(i)$$

The value that $f(i)$ converges to as i approaches c
(Even if $f(c)$ is not defined)

Limit Examples

$$\lim_{i \rightarrow \infty} \frac{1}{i} = 0$$

$$\lim_{i \rightarrow \infty} 6 = 6$$

$$\lim_{i \rightarrow \infty} 6 + \frac{1}{i} = 6$$

$$\lim_{i \rightarrow \infty} i = \infty$$

$$\lim_{i \rightarrow \infty} i - i = 0$$