

# CSE 250 Recitation

January 29 - 30: Summations, Linked Lists



# Welcome

## Introductions

- SAs: Who are we?
- Class: Who are you?

## Participation

- Written exercises during recitation
- Not about getting it right...it's about gaining hands on experience\*
- Good opportunity to ask questions
- **Turned in with your name and UBIT for attendance** (see syllabus)

\* we reserve the right to not count participation with no effort

# Summations – General Form

$$\sum_{i=j}^k f(i) = f(j) + f(j+1) + \dots + f(k)$$

# Summations - Examples

k is 7

f(i) is 5i

$$\sum_{i=4}^7 5i = 20 + 25 + 30 + 35$$

j is 4

f(4)      f(5)      f(6)      f(7)

# Summations - Examples

$n^2$  is  $n^2$        $f(i)$  is  $15n \cdot i$

$$\sum_{i=n}^{n^2} 15n \cdot i = 15n^2 + 15n \cdot (n + 1) + \dots + 15n^3$$

$j$  is  $n$

$f(n)$        $f(n+1)$        $f(n^2)$

The bounds of our summation  
can be unknowns!

# Simplifying the Summation

$$\sum_{i=1}^{n^2} 15n \cdot i$$

Identify the parts of this summation  
are **constant with respect to the  
summation variable**

# Simplifying the Summation

$$\sum_{i=1}^{n^2} 15n \cdot i$$

Identify the parts of this summation are **constant with respect to the summation variable**

(notice how on the previous slide each term had a  $15n$  that did not change...only the value of  $i$  did)

# Simplifying the Summation

$$\sum_{i=n}^{n^2} 15n \cdot i$$

Which of S1, S8, or S9 does this most resemble?

$$S1. \sum_{i=j}^k c = \dots$$

$$S8. \sum_{i=1}^k i = \dots$$

$$S9. \sum_{i=0}^k 2^i = \dots$$

# Simplifying the Summation

$$\sum_{i=1}^{n^2} 15n \cdot i$$

Which of S1, S8, or S9 does this most resemble?

S1.  $\sum_{i=j}^k c = \dots$

S8.  $\sum_{i=1}^k i = \dots$

S9.  $\sum_{i=0}^k 2^i = \dots$

# Simplifying the Summation

$$\sum_{i=n}^{n^2} 15n \cdot i$$

What parts of the summation don't match the rule?

$$\text{S8. } \sum_{i=1}^k i = \dots$$

# Simplifying the Summation

$$\sum_{i=n}^{n^2} 15n \cdot i$$

What parts of the summation don't match the rule?

$$S8. \sum_{i=1}^k i = \dots$$

Notice how the lower bound in S8 is NOT an unknown...it must be exactly 1 to match

# Recitation Exercise

Simplify the summation to its closed form solution using rules on the next slide

$$\sum_{i=n}^{n^2} 15n \cdot i$$

# Summation Rules

$$S1. \sum_{i=j}^k c = (k - j + 1)c$$

$$S2. \sum_{i=j}^k (cf(i)) = c \sum_{i=j}^k f(i)$$

$$S3. \sum_{i=j}^k (f(i) + g(i)) = \left( \sum_{i=j}^k f(i) \right) + \left( \sum_{i=j}^k g(i) \right)$$

$$S4. \sum_{i=j}^k (f(i)) = \left( \sum_{i=\ell}^k (f(i)) \right) - \left( \sum_{i=\ell}^{j-1} (f(i)) \right) \quad (\text{for any } \ell < j)$$

$$S5. \sum_{i=j}^k f(i) = f(j) + f(j+1) + \dots + f(k-1) + f(k)$$

$$S6. \sum_{i=j}^k f(i) = f(j) + \dots + f(\ell-1) + \left( \sum_{i=\ell}^k f(i) \right) \quad (\text{for any } j < \ell \leq k)$$

$$S7. \sum_{i=j}^k f(i) = \left( \sum_{i=j}^{\ell} f(i) \right) + f(\ell+1) + \dots + f(k) \quad (\text{for any } j \leq \ell < k)$$

$$S8. \sum_{i=1}^k i = \frac{k(k+1)}{2}$$

$$S9. \sum_{i=0}^k 2^i = 2^{k+1} - 1$$


$$\sum_{i=n}^{n^2} 15n \cdot i$$

"Bonus" Question


$$\sum_{x=0}^n \sum_{y=1}^k (x + 1)$$

# Linked Lists vs. Arrays

List



Array



# Recitation Exercise

Write out **pseudocode** for the following two algorithms:

1. Find the value of a linked list node at a given index
2. Find the index of a linked list node with a given value (return -1 if the value does not exist)

**Be Precise!**

**If you write good pseudocode, translating it into Java (or C, or Python, or Scala, etc) should be straightforward!**

**Note: PA1 is about LinkedLists!**